



# Mississippi

## Live births, abortions, and pregnancies


Live birth rates, 1982–1992



Live birth rates,\* 1991



Pregnancy rates, 1990



National live birth rate ranking, 1990

Age group		
15-17	18-19	15-19
2nd highest	7th highest	2nd highest


\*Missing bar indicates fewer than 10,000 females or fewer than 20 births. This graph refers to females aged 15–19 years.




# Sexually transmitted diseases

Mississippi

## Gonorrhea rates, 1982–1992



## Gonorrhea rates, 1992



## National ranking,\* 1992

Males – 5th highest    Females – 3rd highest



\*National ranking refers to males and females aged 10–24 years.



## Chlamydia cases and rates, 1988–1992

*Chlamydia data are not available for Mississippi.*

## Acquired immunodeficiency syndrome (AIDS) cumulative cases,\* 1981–1992




\*Missing bar indicates fewer than six cumulative cases.



# Related risk behaviors,\* 1993

Mississippi


Had sexual intercourse (lifetime)



Had sexual intercourse with four or more people (lifetime)




Had sexual intercourse during the past three months (currently sexually active)



Used a condom during last sexual intercourse (among currently sexually active)



Used birth control pills during last sexual intercourse (among currently sexually active)



Injected illegal drugs (lifetime)



\*Related risk behaviors refer to high school students.

Sources: CDC—NCHS and NCCDPHP (birth, abortion, and pregnancy data), NCPS (gonorrhea data); NCID (AIDS data), 1993 Mississippi Youth Risk Behavior Survey (risk behavior data).

